

Massless arbitrary spin fields in AdS_5

R.R. Metsaev

*Department of Theoretical Physics, P.N. Lebedev Physical Institute,
Leninsky prospect 53, 119991, Moscow, Russia*

Abstract

Free arbitrary spin massless and self-dual massive fields propagating in AdS_5 are investigated. We study totally symmetric and mixed symmetry fields on an equal footing. Light-cone gauge action for such fields is constructed. As an example of application of light-cone formalism we discuss AdS/CFT correspondence for massless arbitrary spin AdS fields and corresponding boundary operators at the level of two point function.

Introduction. A study of higher spin gauge theory has two main motivations (see e.g. [1, 2]): Firstly to overcome the well-known barrier of $N \leq 8$ in $d = 4$ supergravity models and, secondly, to investigate if there is a most symmetric phase of superstring theory that leads to the usual string theory as a result of a certain spontaneous breakdown of higher spin symmetries. Another motivation came recently from conjectured duality of free large N conformal $\mathcal{N} = 4$ SYM theory and a theory of massless higher spin fields in AdS_5 [3, 4]. Discussion of this theme in the context of various limits in AdS superstring may be found in [5, 6]. In both the tensionless superstring theory and massless higher spin fields theory all types of massless fields appear in general. This implies that all types of massless fields should be studied. This is what we are doing in this paper.

Let us first formulate the main problem we solve in this letter. Fields propagating in AdS_5 space are associated with positive-energy unitary lowest weight representations of $SO(4, 2)$ group. A positive-energy lowest weight irreducible representation of the $SO(4, 2)$ group denoted as $D(E_0, \mathbf{h})$, is defined by E_0 , the lowest eigenvalue of the energy operator, and by $\mathbf{h} = (h_1, h_2)$, $h_1 \geq |h_2|$, which is the highest weight of the unitary representation of the $SO(4)$ group. The highest weights h_1, h_2 are integers and half-integers for bosonic and fermionic fields respectively. In this paper we investigate the fields whose E_0 and $\mathbf{h}$ are given by

$$E_0 = h_1 + 2, \quad h_1 > |h_2|, \quad (1)$$

$$E_0 > h_1 + 1, \quad h_1 = |h_2|, \quad |h_2| > 1/2. \quad (2)$$

The fields in (1) are massless fields while the ones in (2) are massive self-dual fields. The massless fields with $h_1 \geq 1$, $h_2 = 0, \pm 1/2$ are referred to as totally symmetric fields while both the massless and massive fields with $|h_2| > 1/2$ we refer to as mixed symmetry fields. In manifestly Lorentz covariant formulation the totally symmetric and mixed symmetry fields are the tensor fields whose $SO(4, 1)$ space-time indices have the structure of the respective Young tableaux with one and two rows¹. Actions for the massless totally symmetric integer and half-integer spin fields in AdS_d space were found in [7, 8]. The gauge invariant equations of motion for all types of massless fields (totally and mixed symmetric ones) in AdS_d space for even d have been found in [9, 10]². Massless mixed symmetry AdS_5 fields with arbitrary $\mathbf{h}$ and self-dual massive fields with arbitrary E_0 have not been described at the field theoretical level so far³. In this paper we develop a light-cone gauge formulation for such fields at the action level. Using the light-cone gauge formalism in AdS space developed in [14], we describe both the totally and mixed symmetry fields on an equal footing. Since, by analogy with flat space, we expect that a quantization of the Green-Schwarz AdS superstring with Ramond-Ramond charge will be

¹We note that the case $\mathbf{h} = (1, 0)$ corresponds to spin one Maxwell field, the case $\mathbf{h} = (2, 0)$ is the graviton, and the cases $\mathbf{h} = (1, \pm 1)$ correspond to 2-form potentials.

²The particular case of massless field corresponding to $\mathbf{h} = (2, 1)$, for $d \geq 6$ was described at the action level in [11]. Gauge invariant actions for AdS_4 massless fields have been established in [12, 13]. Various gauge invariant descriptions of totally symmetric massless fields in terms of the potentials in AdS_d may be found in [14, 15, 16]. Massive self-dual arbitrary spin fields in AdS_3 were investigated in [17]. Discussion of massive totally symmetric fields in AdS_d , $d \geq 4$, may be found in [18, 19].

³Group theoretical description of representation of $so(4, 2)$ algebra for arbitrary $\mathbf{h}$ and discrete values of E_0 via oscillator method [20] may be found, e.g., in [21]. Lorentz covariant equations of motion for AdS_5 self-dual massive fields with special values of E_0 were discussed in [22].

available only in the light-cone gauge [23, 24] it seems that from the stringy perspective of *AdS/CFT* correspondence the light-cone approach is the fruitful direction to go.

Light-cone gauge action. Let $\phi(x)$ be a arbitrary spin field propagating in AdS_5 space. If we collect spin degrees of freedom in a generating function $|\phi(x)\rangle$ then a light-cone gauge action for this field can be cast into the following ‘covariant form’[14]⁴

$$S_{l.c.} = \frac{1}{2} \int d^5x \langle \phi(x) | (\square - \frac{1}{z^2} A) | \phi(x) \rangle, \quad \square = 2\partial^+ \partial^- + \partial_i^2 + \partial_z^2, \quad (3)$$

where an A is some operator not depending on space-time coordinates and their derivatives. This operator acts only on spin indices of fields collected in ket-vector $|\phi\rangle$. We call the operator A the *AdS* mass operator. In the remainder of this paper we find it convenient to Fourier transform to momentum space for all coordinates except for the radial z and the light-cone time x^+ :

$$|\phi(x^+, x^-, x^i, z)\rangle = \int \frac{d^2p d\beta}{(2\pi)^{3/2}} e^{i(\beta x^- + p^i x^i)} |\phi(x^+, \beta, p^i, z)\rangle, \quad (4)$$

where we use β to denote a momentum in light-cone direction: $\beta \equiv p^+$. Now we rewrite the action in the Hamiltonian form

$$S_{l.c.} = \int dx^+ d^2p d\beta dz \langle \phi(x^+, -p, z) | \beta (i\partial^- + P^-) | \phi(x^+, p, z) \rangle, \quad (5)$$

where P^- is the (minus) Hamiltonian density

$$P^- = \frac{-p_i^2 + \partial_z^2}{2\beta} - \frac{1}{2z^2\beta} A. \quad (6)$$

In Eq. (5) and below a momentum p as argument of wave function $\phi(x^+, p, z)$ designates the set $\{p^i, \beta\}$. All that remains to fix action explicitly is to find the *AdS* mass operator. The explicit form of this operator depends on what kind of realization of spin degrees of freedom we use. Let us first consider the bosonic fields.

Bosonic fields. To describe light-cone gauge formulation of the $D(E_0, \mathbf{h})$ massless or self-dual massive bosonic field $|\phi\rangle$ we introduce a *complex-valued* totally symmetric traceless $so(3)$ tensor field $\phi^{I_1 \dots I_{h_1}}$. As usual to avoid cumbersome tensor expressions we introduce creation and annihilation oscillators α^I and $\bar{\alpha}^I$

$$[\bar{\alpha}^I, \alpha^J] = \delta^{IJ}, \quad \bar{\alpha}^I |0\rangle = 0, \quad (7)$$

and the generating function $|\phi\rangle$ defined by

$$|\phi(x^+, p, z)\rangle \equiv \alpha^{I_1} \dots \alpha^{I_{h_1}} \phi^{I_1 \dots I_{h_1}}(x^+, p, z) |0\rangle. \quad (8)$$

The *AdS* mass operator is found then to be

⁴ We use parametrization of AdS_5 space in which $ds^2 = (-dx_0^2 + dx_i^2 + dx_3^2 + dz^2)/z^2$. Light-cone coordinates in $\pm$ directions are defined as $x^\pm = (x^3 \pm x^0)/\sqrt{2}$ and x^+ is taken to be a light-cone time. Unless otherwise specified, we adopt the conventions: $i, j = 1, 2$. $\partial_i \equiv \partial/\partial x^i$, $\partial_z \equiv \partial/\partial z$, $\partial^\pm = \partial_\mp \equiv \partial/\partial x^\mp$. We use indices I, J for $so(3)$ tangent space vectors $X^I = (X^i, X^z)$. Sometimes, instead of $so(2)$ vector indices $i, j = 1, 2$ we use complex frame indices $i, j = x, \bar{x}$. In this notation a $so(2)$ vector X^i is representable by $X^x, X^{\bar{x}}$ and for scalar product we have the decomposition $X^i Y^i = X^x Y^{\bar{x}} + X^{\bar{x}} Y^x$.

$$A = -\frac{1}{2}m^{ij}m^{ij} - \frac{1}{4}, \quad m^{ij} \equiv M^{ij} - i\epsilon^{ij}c, \quad (9)$$

$\epsilon^{12} = -\epsilon^{21} = 1$, where the $so(2)$ spin operator M^{ij} takes the standard form

$$M^{ij} = \alpha^i \bar{\alpha}^j - \alpha^j \bar{\alpha}^i, \quad (10)$$

and the number c is given by

$$c = \begin{cases} h_2, & \text{for massless fields (1),} \\ (E_0 - 2)\text{sign } h_2, & \text{for massive self-dual fields (2),} \end{cases} \quad (11)$$

where $\text{sign } h = +1(-1)$ for $h > 0(h < 0)$. Details of derivation may be found in Appendix B. A few comments are in order.

i) The light-cone gauge action for the mixed symmetry $D(E_0, \mathbf{h})$ field, ($|h_2| > 0$), is formulated in terms of the *complex-valued* rank h_1 totally symmetric traceless $so(3)$ tensor field. Number of physical degrees of freedom (DoF) of such field is equal to $2(2h_1 + 1)$, i.e., for the field in (1), it depends only on h_1 and does not on h_2 .

(ii) For the case of massless totally symmetric field ($h_2 = 0$) we can impose reality condition on the complex-valued $so(3)$ tensor field and this gives the standard light-cone description of the massless totally symmetric field [14]. Number of physical DoF of such field is equal to $2h_1 + 1$.

(iii) The h_2 of AdS_5 fields is similar to helicity of massless fields in flat four dimensional space. As is well known number of physical DoF of the latter fields does not depend on helicity and is equal to two. If AdS_5 field $|\phi\rangle$ has helicity h_2 then its complex-conjugated $\langle\phi|$ has opposite helicity $-h_2$, i.e., the light-cone gauge action (5) is formulated in terms of fields having opposite values of h_2 .

Fermionic fields. In this case h_1 and h_2 are half-integers and in what follows we use notation $h'_1 = h_1 - \frac{1}{2}$. To discuss light-cone gauge formulation of $D(E_0, \mathbf{h})$ fermionic field we introduce complex-valued traceless $so(3)$ spin-tensor field $\phi_\gamma^{I_1 \dots I_{h'_1}}$, $\gamma = 1, 2$, which is symmetric in $I_1, \dots, I_{h'_1}$. In addition to the bosonic creation and annihilation oscillators (7) we introduce fermionic oscillators $\eta^\gamma, \bar{\eta}^\gamma$

$$\{\bar{\eta}^\alpha, \eta^\gamma\} = \delta^{\alpha\gamma}, \quad \bar{\eta}^\gamma|0\rangle = 0. \quad (12)$$

For this case the Fock vector $|\phi\rangle$ defined by

$$|\phi(x^+, p, z)\rangle \equiv \alpha^{I_1} \dots \alpha^{I_{h'_1}} \eta^\gamma \phi_\gamma^{I_1 \dots I_{h'_1}}(x^+, p, z)|0\rangle \quad (13)$$

is assumed to satisfy the constraint

$$(\bar{\alpha}^i \sigma^i - \bar{\alpha}^z) \sigma^3 \bar{\eta}|\phi\rangle = 0, \quad (14)$$

where σ^i, σ^3 are Pauli matrices. This constraint amounts to requiring the spin-tensor field $\phi_\alpha^{I_1 \dots I_{h'_1}}$ transforms in irreducible spin h_1 representation of $so(3)$ algebra. The operator A and the number c take then the form given by formulas (9),(11) in which the spin operator M^{ij} (10) should be replaced by

$$M^{ij} = \alpha^i \bar{\alpha}^j - \alpha^j \bar{\alpha}^i + \frac{i}{2} \epsilon^{ij} \eta \sigma^3 \bar{\eta}. \quad (15)$$

As a side of remark we note that these results can be presented in an equivalent form which is somewhat convenient for evaluation. Instead of tensor fields (8),(13) we can introduce $2h_1 + 1$ complex-valued fields ϕ_m which with the help of a complex-valued variable ζ we collect in the $|\phi\rangle$ as

$$|\phi(x^+, p, z)\rangle = \sum_{m=-h_1}^{h_1} \zeta^{h_1-m} \phi_m(x^+, p, z), \quad (16)$$

where for bosons $m = 0, \pm 1, \dots, \pm h_1$, while for fermions $m = \pm 1/2, \pm 3/2, \dots, \pm h_1$. The ζ , which we refer to as projective variable, is used to formulate results entirely in terms of the generating function $|\phi\rangle$. The scalar product for such generating functions is defined to be

$$\langle \phi | \varphi \rangle = \int \frac{d\zeta d\bar{\zeta}}{(1 + \frac{1}{2}\zeta\bar{\zeta})^{2h_1+2}} \overline{\phi(\zeta)} \varphi(\zeta). \quad (17)$$

The *AdS* mass operator for this realization takes the form

$$A = (\zeta \partial_\zeta + a)^2 - \frac{1}{4}, \quad a = c - h_1, \quad (18)$$

where the number c is given by (11). In the complex notation the spin operator M^{ij} is representable by $M^{x\bar{x}}$ which takes the form

$$M^{x\bar{x}} = -\zeta \partial_\zeta + h_1, \quad \partial_\zeta \equiv \partial/\partial\zeta. \quad (19)$$

We note that the representation for the operators A (18) and $M^{x\bar{x}}$ (19) is valid for both the integer and half-integer fields, i.e., while using the projective variable ζ it is not necessary to introduce anticommuting oscillators (12) to discuss the half-integer fields.

Global symmetries of light-cone gauge action. We turn now to discussion of global $so(4, 2)$ symmetries of the light-cone gauge action. The choice of the light-cone gauge spoils the manifest global symmetries, and in order to demonstrate that these global invariances are still present one needs to find the Noether charges which generate them⁵. The field theoretical representation for Noether charges in *AdS* light-cone gauge formalism takes the standard form [14]

$$\hat{G} = \int dz d^2 p d\beta \beta \langle \phi(x^+, -p, z) | G | \phi(x^+, p, z) \rangle, \quad (20)$$

where G are differential operators acting on physical field $|\phi\rangle$. Noether charges (generators) can be split into two groups: kinematical and dynamical generators (see Appendix A). Representation of the kinematical generators is given by

$$P^i = p^i, \quad P^+ = \beta, \quad (21)$$

$$D = ix^+ P^- - \partial_\beta \beta - \partial_{p^i} p^i + z \partial_z + \frac{3}{2}, \quad (22)$$

$$J^{+-} = ix^+ P^- + \partial_\beta \beta, \quad (23)$$

$$J^{+i} = ix^+ p^i + \partial_{p^i} \beta, \quad (24)$$

⁵These charges play a crucial role in formulating interaction vertices in field theory. Application of Noether charges in formulating superstring field theories may be found in [25],[26]

$$J^{ij} = p^i \partial_{p^j} - p^j \partial_{p^i} + M^{ij}, \quad (25)$$

$$K^+ = \frac{1}{2}(2ix^+ \partial_\beta - \partial_{p^i}^2 + z^2)\beta + ix^+ D, \quad (26)$$

$$K^i = \frac{1}{2}(2ix^+ \partial_\beta - \partial_{p^j}^2 + z^2)p^i - \partial_{p^i} D - M^{ij} \partial_{p^j} - M^{zi} z + iM^{i-} x^+, \quad (27)$$

where $\partial_{p^i} \equiv \frac{\partial}{\partial p^i}$, $\partial_\beta \equiv \frac{\partial}{\partial \beta}$. The dynamical generator P^- is given by (6) while the remaining ones are

$$J^{-i} = -\partial_\beta p^i + \partial_{p^i} P^- + M^{-i}, \quad (28)$$

$$K^- = \frac{1}{2}(2ix^+ \partial_\beta - \partial_{p^i}^2 + z^2)P^- - \partial_\beta D + \frac{1}{\beta} p^i \partial_{p^j} M^{ij} + \frac{1}{\beta} (zp^i - \partial_z \partial_{p^i}) M^{zi} - \frac{\partial_{p^i}}{2z\beta} [M^{zi}, A] + \frac{1}{\beta} B, \quad (29)$$

where ‘nonlinear’ spin operator $M^{-i} = -M^{i-}$ is given by

$$M^{-i} \equiv \frac{1}{\beta} \left(M^{ij} p^j + M^{zi} \partial_z + \frac{1}{2z} [M^{zi}, A] \right). \quad (30)$$

The operators B , M^{zi} are acting only on spin degrees of freedom and are given by

$$B = -M^{zi} M^{zi} + \frac{i}{2} c \epsilon^{ij} M^{ij}, \quad (31)$$

$$M^{zi} = \alpha^z \bar{\alpha}^i + \alpha^i \bar{\alpha}^z + \frac{1}{2} \eta \sigma^i \bar{\eta}. \quad (32)$$

For the case of bosonic fields we should simply ignore the fermionic term in expression for M^{zi} (32). We note that the spin operators M^{ij} , M^{zi} satisfy commutation relations of $so(3)$ algebra

$$[M^{zi}, M^{zj}] = M^{ij}, \quad [M^{ij}, M^{zk}] = \delta^{jk} M^{zi} - \delta^{ik} M^{zj}, \quad (33)$$

and a realization of the spin operator M^{zi} in terms of projective variable ζ takes the form

$$M^{zx} = \partial_\zeta, \quad M^{z\bar{x}} = -\frac{1}{2} \zeta^2 \partial_\zeta + h_1 \zeta. \quad (34)$$

Since the light-cone gauge action (5) is invariant with respect to the global symmetries generated by $so(4, 2)$ algebra, $\delta_{\hat{G}}|\phi\rangle = G|\phi\rangle$, the formalism we discuss is sometimes referred to as an off shell light-cone formulation [25].

Ground state of AdS_5 fields. In this section we would like to demonstrate that the massless and self-dual massive fields discussed above are indeed carriers of the positive-energy unitary lowest weight representations of $SO(4, 2)$ group labeled by $D(E_0, \mathbf{h})$ where E_0 is given by Eqs. (1),(2). To do this we evaluate ground state of AdS fields. All that requires is to demonstrate that E_0 for ground states of massless and self-dual fields satisfy the basic relations (1),(2). To this end we begin with definition of ground state $|v\rangle$ (vacuum). The most convenient way to do this is to use $so(4, 2)$ algebra taken to be in energy basis (see Appendix A). In this basis one has energy operator $J^{w\bar{w}}$, deboost

operators $J^{wy}, J^{w\bar{y}}, J^{wi}$, boost operators, $J^{\bar{w}y}, J^{\bar{w}\bar{y}}, J^{\bar{w}i}$ and six operators, $J^{iy}, J^{i\bar{y}}, J^{y\bar{y}}, J^{ij}$, which are generators of the $so(4)$ algebra⁶. The vacuum forms a linear space which is invariant under the action of the energy operator $J^{w\bar{w}}$ and generators of the $so(4)$ algebra. In other words the $|v\rangle$ is i) the eigenvalue vector of $J^{w\bar{w}}$; ii) a weight $\mathbf{h}$ representation of the $so(4)$ algebra. Now the vacuum $|v\rangle$ is defined by the relations

$$J^{w\bar{w}}|v\rangle = E_0|v\rangle, \quad (35)$$

$$J^{wy}|v\rangle = 0, \quad J^{w\bar{y}}|v\rangle = 0, \quad (36)$$

$$J^{wi}|v\rangle = 0. \quad (37)$$

Eq. (35) reflects the fact that the $|v\rangle$ is the eigenvector of the energy operator $J^{w\bar{w}}$, while Eqs. (36),(37) tell us that E_0 is the lowest value of the energy: since the deboost operators decrease the energy they should evidently annihilate the $|v\rangle$. Just as in the ordinary Poincaré quantum field theory one can build the representation Fock space $D(E_0, \mathbf{h})$ by acting with boost operators $J^{\bar{w}y}, J^{\bar{w}\bar{y}}, J^{\bar{w}i}$ on the vacuum states $|v\rangle$.

Without loss of generality and to simplify our presentation we i) evaluate representative of ground state denoted by $|v^{E_0, \mathbf{h}}\rangle$ which is a highest weight vector of representation of the $so(4)$ algebra; ii) evaluate ground state for $x^+ = 0$ and $\beta > 0$; iii) drop the $|v^{E_0, \mathbf{h}}\rangle$ in equations and instead of equations like $X|v^{E_0, \mathbf{h}}\rangle = Y|v^{E_0, \mathbf{h}}\rangle$ we write simply $X \approx Y$. In addition to (35)-(37) the $|v^{E_0, \mathbf{h}}\rangle$ satisfies, by definition, the following constraints

$$J^{y\bar{y}} \approx h_1, \quad J^{x\bar{x}} \approx h_2, \quad (38)$$

$$J^{yi} \approx 0. \quad (39)$$

Next step is to rewrite Eqs. (35)-(39) in terms of generators in light-cone basis (21)-(29). Making use of (37),(39) and formulas (66),(68) we get the equations

$$P^i + J^{+i} \approx 0, \quad (40)$$

$$K^i - J^{-i} \approx 0. \quad (41)$$

Using Eqs. (36) and formulas (62),(63) we can express P^- and K^- in terms of P^+, K^+, D and J^{+-} . Exploiting this representation for P^-, K^- in (60),(61) we get from (35) and the 1st equation in (38) the following equations

$$P^+ + \frac{1}{2}(-D + J^{+-}) \approx \frac{1}{2}(E_0 + h_1), \quad (42)$$

$$K^+ + \frac{1}{2}(D + J^{+-}) \approx \frac{1}{2}(E_0 - h_1). \quad (43)$$

Now making use of concrete representation for generators in terms of differential operators (21)-(29) we can find ground state explicitly. Using Eqs. (40),(42),(43) we find respective dependence of the ground state $|v^{E_0, \mathbf{h}}\rangle$ on momenta p^i , β and radial coordinate z :

$$|v^{E_0, \mathbf{h}}(p, \beta, z)\rangle = \exp\left(-\frac{1}{2\beta}p^i p^i - \frac{1}{2}\beta z^2 - \beta z^{E_0 - h_1 - \frac{3}{2}}\beta^{E_0 - 1}|v_0\rangle\right), \quad (44)$$

⁶In complex notation J^{ij} is representable by $J^{x\bar{x}}$ while $so(2)$ vectors like J^{wi} are representable by J^{wx} and $J^{w\bar{x}}$

where $|v_0\rangle$ depends only on ζ . To find a dependence of $|v_0\rangle$ on ζ we use the 2nd equation in (38) which with the help of (19) gives

$$|v_0\rangle = \zeta^{h_1-h_2}. \quad (45)$$

Formulas (44),(45) give explicit representation for the ground state which we use to find desired constraint on E_0 (1),(2). To this end we analyze Eqs. (41) which for $i = x$ and $i = \bar{x}$ lead to the equations

$$(E_0 - h_1 - 1 + a + \zeta \partial_\zeta) M^{zx} |v_0\rangle = 0, \quad (46)$$

$$(E_0 - h_1 - 1 - a - \zeta \partial_\zeta) M^{z\bar{x}} |v_0\rangle = 0. \quad (47)$$

From now on we analyze the cases $h_1 > |h_2|$ and $h_1 = |h_2|$ in turn.

i) If $h_1 > |h_2|$ then Eqs. (46),(47) give

$$E_0 - 2 + a - h_2 = 0, \quad E_0 - 2h_1 - 2 - a + h_2 = 0. \quad (48)$$

It is easy to see that solution to these equations is $E_0 = h_1 + 2$, $a = h_2 - h_1$. These E_0 and a correspond to massless fields (see (1),(11),(18)).

ii) If $h_1 = h_2$, $h_2 > 0$ then Eq. (46) is satisfied automatically while Eq. (47) gives $a = E_0 - 2 - h_1$.

iii) If $h_1 = -h_2$, $h_2 < 0$ then Eq. (47) is satisfied automatically while Eq. (46) gives $a = 2 - E_0 - h_1$. Thus the cases ii) and iii) lead to values of a corresponding to the self-dual massive fields (see (18),(11)). As to restriction on E_0 (2) it follows simply from the requirement that the ground state (44) be square-integrable with respect to z : $\int dz ||v^{E_0, h}\rangle|^2 < \infty$. A requirement of square-integrability with respect to remaining variables does not lead to new constraints.

AdS/CFT correspondence. In spite of its Lorentz noncovariance, a light-cone formalism offers conceptual and technical simplification of approaches to various problems of modern quantum field theory. One of problems which allows to demonstrate efficiency of a light-cone approach is *AdS/CFT* correspondence [27, 28, 29]. In the Lorentzian signature *AdS/CFT* correspondence at the level of state/operator matching between the bulk massless arbitrary spin totally symmetric fields and the corresponding boundary operators was demonstrated in [14]. Here we would like to demonstrate this correspondence in the Euclidean signature at the level of two point function⁷. In light-cone gauge the massless arbitrary spin field in AdS_d is described by an totally symmetric traceless $so(d-2)$ tensor field $\phi^{I_1 \dots I_s}$, $I = 1, \dots, d-2$. This tensor can be decomposed into traceless totally symmetric tensors of $so(d-3)$ algebra $\phi^{I_1 \dots I_s} = \sum_{s'=0}^s \phi_{s'}^{i_1 \dots i_{s'}}$, $i = 1, \dots, d-3$. The Euclidean light-cone gauge action⁸ takes then the following simple form [14, 32]

$$S_{l.c.}^E = \sum_{s'=0}^s \frac{1}{2} \int d^d x (|d\phi_{s'}|^2 + \frac{1}{z^2} A_{s'} |\phi_{s'}|^2), \quad (49)$$

where an eigenvalue of the *AdS* mass operator for $\phi^{i_1 \dots i_{s'}}$ is given by

⁷Discussion of *AdS/CFT* correspondence for spin one Maxwell field, $s = 1$, and graviton, $s = 2$, may be found in [29] and [30, 31] respectively.

⁸Note that only in this section we use the Euclidean signature. For $d = 4$ the *AdS* mass operator is equal to zero for all massless fields. Here we restrict our attention to the dimensions $d \geq 5$.

$$A_{s'} = \nu_{s'}^2 - \frac{1}{4}, \quad \nu_{s'} \equiv s' + \frac{d-5}{2}. \quad (50)$$

Attractive feature of this action is that there are no contractions of tensor indices of fields with the ones of space derivatives, i.e., the action looks like a sum of actions for ‘scalar’ fields with different mass terms. This allows us to extend the analysis of Ref.[29] in a rather straightforward way. Using Green’s function method a solution to equations of motion

$$\left(-\partial_{\mathbf{x}}^2 - \partial_z^2 + \frac{1}{z^2}(\nu_{s'}^2 - \frac{1}{4})\right)\phi^{i_1 \dots i_{s'}}(\mathbf{x}, z) = 0, \quad \mathbf{x} \equiv (x^1, \dots, x^{d-1}) \quad (51)$$

is found to be

$$\phi^{i_1 \dots i_{s'}}(\mathbf{x}, z) = \int d\mathbf{x}' \frac{z^{\nu_{s'} + \frac{1}{2}}}{(z^2 + |\mathbf{x} - \mathbf{x}'|^2)^{\nu_{s'} + \frac{d-1}{2}}} \tilde{\mathcal{O}}^{i_1 \dots i_{s'}}(\mathbf{x}'). \quad (52)$$

As was expected [14] this solution behaves for $z \rightarrow 0$ like $z^{-\nu_{s'} + \frac{1}{2}} \tilde{\mathcal{O}}(\mathbf{x})$. Plugging this solution into the action (49) and evaluating a surface integral gives

$$S_{l.c.}^E = \sum_{s'=0}^s \frac{1}{2}(\nu_{s'} + \frac{1}{2}) \int d\mathbf{x} d\mathbf{x}' \frac{\tilde{\mathcal{O}}^{i_1 \dots i_{s'}}(\mathbf{x}) \tilde{\mathcal{O}}^{i_1 \dots i_{s'}}(\mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|^{2\nu_{s'} + d-1}} \quad (53)$$

This is $so(d-3)$ decomposition of two point function of spin s conserved current.

For the case of AdS_5 this analysis can be generalized to arbitrary $D(E_0, h_1, h_2)$ fields (1),(2) in a rather straightforward way. The most convenient way to do this is to use representation for AdS mass operator given in (18). All that is required is to replace in above formulas the $\nu_{s'}$ by $\hat{\nu} = |\zeta \partial_\zeta + a|$, and the fields $\phi_{s'}$ by $|\phi\rangle$ (16). Repetition of the procedure outlined above gives

$$S_{l.c.}^E = \frac{1}{2} \int d\mathbf{x} d\mathbf{x}' \langle \tilde{\mathcal{O}}(\mathbf{x}) | \frac{\hat{\nu} + \frac{1}{2}}{|\mathbf{x} - \mathbf{x}'|^{2\hat{\nu} + 4}} | \tilde{\mathcal{O}}(\mathbf{x}') \rangle. \quad (54)$$

As in (16) the ket-notation $|\mathcal{O}\rangle$ is used to indicate dependence of conformal operator on the complex-valued variable ζ . A scalar product $\langle \mathcal{O} | \mathcal{O}' \rangle$ is defined as in (17).

Acknowledgments. This work was supported by the INTAS project 991590, by the RFBR Grant No.02-02-17067, and RFBR Grant for Leading Scientific Schools, Grant No. 01-02-30024.

Appendix A. Generic commutation relations of $so(4,2)$ algebra generators J^{CF} , $C, F = 0', 0, 1, \dots, 4$, we use are

$$[J^{AB}, J^{CF}] = \eta^{BC} J^{AF} + 3 \text{ terms}, \quad \eta^{CF} = (-, -, +, \dots, +). \quad (55)$$

In the conformal algebra notation the J^{CF} split into translations P^a , conformal boosts K^a , dilatation D , and $so(3,1)$ rotations J^{ab} , $a, b = 0, 1, 2, 3$. Commutation relations in this basis are

$$[D, P^a] = -P^a, \quad [D, K^a] = K^a, \quad [P^a, K^b] = \eta^{ab} D - J^{ab}, \quad (56)$$

$$[P^a, J^{bc}] = \eta^{ab} P^c - \eta^{ac} P^b, \quad [K^a, J^{bc}] = \eta^{ab} K^c - \eta^{ac} K^b, \quad (57)$$

$$[J^{ab}, J^{ce}] = \eta^{bc} J^{ae} + 3 \text{ terms}, \quad \eta^{ab} = (-, +, +, +). \quad (58)$$

In the light-cone basis these generators split into two groups:

$$P^+, P^i, K^+, K^i, J^{+i}, J^{+-}, J^{ij}, D, \quad (59)$$

which we refer to as kinematical generators, and P^-, K^-, J^{-i} which we refer to as dynamical generators. The kinematical generators have positive or zero J^{+-} charges, while dynamical generators have negative J^{+-} charges. Commutation relations in light-cone basis are obtainable from (56)-(58) with the light-cone metric having the following non-vanishing elements $\eta^{+-} = \eta^{-+} = 1$, $\eta^{ij} = \delta^{ij}$.

In energy basis the generators J^{CF} split into energy operator $J^{w\bar{w}}$, deboost operators $J^{wy}, J^{w\bar{y}}, J^{wi}$, boost operators $J^{\bar{w}y}, J^{\bar{w}\bar{y}}, J^{\bar{w}i}$, and $so(4)$ algebra generators $J^{iy}, J^{i\bar{y}}, J^{y\bar{y}}, J^{ij}$. Commutation relations in this basis are obtainable from (55) with the metric having the following non-vanishing elements $\eta^{x\bar{x}} = \eta^{y\bar{y}} = -\eta^{w\bar{w}} = 1$ (in the $so(2)$ notation $\eta^{ij} = \delta^{ij}$). Inter-relation of the generators in light-cone and energy bases is given by

$$J^{w\bar{w}} = \frac{1}{2}(P^+ - P^- + K^+ - K^-), \quad (60)$$

$$J^{y\bar{y}} = \frac{1}{2}(P^+ + P^- - K^+ - K^-), \quad (61)$$

$$J^{wy} = \frac{1}{2}(P^- + K^+ + D + J^{+-}), \quad (62)$$

$$J^{w\bar{y}} = \frac{1}{2}(P^+ + K^- - D + J^{+-}), \quad (63)$$

$$J^{\bar{w}y} = \frac{1}{2}(-P^+ - K^- - D + J^{+-}), \quad (64)$$

$$J^{\bar{w}\bar{y}} = \frac{1}{2}(-P^- - K^+ + D + J^{+-}), \quad (65)$$

$$J^{wi} = \frac{1}{2}(P^i + K^i + J^{+i} - J^{-i}), \quad (66)$$

$$J^{\bar{w}i} = \frac{1}{2}(-P^i - K^i + J^{+i} - J^{-i}), \quad (67)$$

$$J^{yi} = \frac{1}{2}(P^i - K^i + J^{+i} + J^{-i}), \quad (68)$$

$$J^{\bar{y}i} = \frac{1}{2}(-P^i + K^i + J^{+i} + J^{-i}). \quad (69)$$

Appendix B. In this appendix we explain a procedure of derivation of the AdS mass operator and the operator B which enter in the definition of the light-cone gauge action and representation of the generators of $so(4, 2)$ algebra on physical fields. As was demonstrated in [14] the operators A and B should satisfy the basic defining equations

$$[A, M^{ij}] = 0, \quad (70)$$

$$2\{M^{zi}, A\} - [[M^{zi}, A], A] = 0, \quad (71)$$

$$-[M^{zi}, [M^{zj}, A]] + \{M^{il}, M^{lj}\} + \{M^{zi}, M^{zj}\} = -2\delta^{ij}B. \quad (72)$$

To find solution of these equations we use representation for spin operators M^{ij}, M^{zi} in terms of projective complex-valued variable ζ given in (19),(34). In general the A is an

operator depending on ζ and ∂_ζ . Equation (70) tells that the A depends only on $\hat{N}$, where $\hat{N} \equiv \zeta \partial_\zeta$. We now are looking for $A = A(\hat{N})$ satisfying the remaining equations (71),(72). Making use of the obvious relations

$$A(\hat{N})\partial_\zeta = \partial_\zeta A(\hat{N} - 1), \quad A(\hat{N})\zeta = \zeta A(\hat{N} + 1), \quad (73)$$

we get from Eqs. (71) for $i = x$ and $i = \bar{x}$ the respective functional equations

$$2(A(\hat{N}) + A(\hat{N} - 1)) - (A(\hat{N}) - A(\hat{N} - 1))^2 = 0, \quad (74)$$

$$2(A(\hat{N}) + A(\hat{N} + 1)) - (A(\hat{N}) - A(\hat{N} + 1))^2 = 0. \quad (75)$$

It is easy to see that solution of these equations does not involve higher than second-order terms in $\hat{N}$. Making use of general anzats $A = q\hat{N}^2 + r\hat{N} + t$ we find that $q = 1$, $r = 2a$, $t = a^2 - 1/4$, where a is arbitrary constant. Thus we get $A = (\hat{N} + a)^2 - 1/4$. To find the operator B we use Eq. (72) for $i = x$, $j = \bar{x}$. Solution to this equation is found to be

$$B = -\{M^{zx}, M^{z\bar{x}}\} - (a + h_1)M^{x\bar{x}}. \quad (76)$$

The remaining unknown number a can be fixed in various ways. One way to fix a is to consider the ground state (see Section below Eq.(34)). Here we would to outline procedure of finding a by using technique of Casimir operators. Let Q be the second order Casimir operator. Making use of expressions for generators (21)-(29) the equation $(Q - \langle Q \rangle)|\phi\rangle = 0$, where $\langle Q \rangle$ is the eigenvalue of the Q in $D(E_0, \mathbf{h})$

$$-\langle Q \rangle = E_0(E_0 - 4) + h_1(h_1 + 2) + h_2^2, \quad (77)$$

can be cast into the form (for details see Section 4 in Ref.[14])

$$-A + 2B + \frac{1}{2}M_{ij}^2 + \frac{15}{4} - \langle Q \rangle = 0. \quad (78)$$

Plugging solution for A and B into (78) we obtain the equation

$$(E_0 - 2)^2 - h_1^2 + h_2^2 - (a + h_1)^2 = 0. \quad (79)$$

In the same way an evaluation of the third order Casimir operator gives the equation $h_1(h_1 + a) = (E_0 - 2)h_2$. For massive fields (2) this equation and (79) lead to a given in (18),(11). For the case of $h_1 > |h_2|$ we get $E_{0\pm} = 2 \pm h_1$ and the physical relevant value E_{0+} (1) can be fixed by unitarity condition (see e.g. [9, 11]). Making use of normalization for Levi-Civita tensor in complex frame $\epsilon^{x\bar{x}} = -i$ it is easy to see that the operators A (18) and B (76) can be cast into $so(2)$ covariant form (9),(31). The most convenient way to relate the spin operators in $so(2)$ covariant form (15),(32) and the ones in (19),(34) is to pass from the oscillators (7) to complex-valued variables a^i , a^z subject to the constraint $a^i a^i - a^z a^z = 0$. For simplicity we consider the bosonic case. In terms of a^i , a^z the $so(2)$ covariant form of the spin operators is

$$M^{ij} = a^i \partial_{a^j} - a^j \partial_{a^i}, \quad M^{zi} = a^z \partial_{a^i} + a^i \partial_{a^z}. \quad (80)$$

Introducing the $\zeta = a^z/a^x$ one can make sure that a representation of the operators (80) on a monomial having h_1 th power in a^i , a^z takes the same form as in (19),(34).

References

- [1] M. A. Vasiliev, Fortsch. Phys. **35**, 741 (1987)
- [2] R. R. Metsaev, “IIB supergravity and various aspects of light-cone formalism in AdS space-time,” arXiv:hep-th/0002008. Talk given at International Workshop on Supersymmetries and Quantum Symmetries, Moscow, Russia, 27-31 Jul 1999.
- [3] P. Haggi-Mani and B. Sundborg, JHEP **0004**, 031 (2000) [arXiv:hep-th/0002189].
- [4] B. Sundborg, Nucl. Phys. Proc. Suppl. **102**, 113 (2001) [arXiv:hep-th/0103247].
- [5] A. A. Tseytlin, arXiv:hep-th/0201112.
- [6] E. Sezgin and P. Sundell, JHEP **0109**, 036 (2001) [arXiv:hep-th/0105001].
- [7] V. E. Lopatin and M. A. Vasiliev, Mod. Phys. Lett. A **3**, 257 (1988).
- [8] M. A. Vasiliev, Nucl. Phys. B **301**, 26 (1988).
- [9] R. R. Metsaev, Phys. Lett. B **354**, 78 (1995); Phys. Lett. B **419**, 49 (1998) [arXiv:hep-th/9802097];
- [10] R. R. Metsaev, “Arbitrary spin massless bosonic fields in d-dimensional anti-de Sitter space,” arXiv:hep-th/9810231. Talk given at International Seminar on Supersymmetries and Quantum Symmetries, Dubna, Russia, 22-26 Jul 1997.
- [11] L. Brink, R. R. Metsaev and M. A. Vasiliev, Nucl. Phys. B **586**, 183 (2000) [arXiv:hep-th/0005136].
- [12] C. Fronsdal, Phys. Rev. D **20**, 848 (1979).
- [13] J. Fang and C. Fronsdal, Phys. Rev. D **22**, 1361 (1980).
- [14] R. R. Metsaev, Nucl. Phys. B **563**, 295 (1999) [arXiv:hep-th/9906217].
- [15] A. Y. Segal, arXiv:hep-th/0103028.
- [16] I. L. Buchbinder, A. Pashnev and M. Tsulaia, Phys. Lett. B **523**, 338 (2001) [arXiv:hep-th/0109067].
- [17] R. R. Metsaev, Nucl. Phys. Proc. Suppl. **102**, 100 (2001) [arXiv:hep-th/0103088].
- [18] S. Deser and A. Waldron, Phys. Rev. Lett. **87**, 031601 (2001) [arXiv:hep-th/0102166]; Nucl. Phys. B **607**, 577 (2001) [arXiv:hep-th/0103198].
- [19] Y. M. Zinoviev, arXiv:hep-th/0108192.
- [20] I. Bars and M. Gunaydin, Commun. Math. Phys. **91**, 31 (1983).
- [21] M. Gunaydin, D. Minic and M. Zagermann, Nucl. Phys. B **534**, 96 (1998) [Erratum-ibid. B **538**, 531 (1998)] [arXiv:hep-th/9806042].
- [22] E. Sezgin and P. Sundell, JHEP **0109**, 025 (2001) [arXiv:hep-th/0107186].
- [23] R. R. Metsaev and A. A. Tseytlin, Phys. Rev. D **63**, 046002 (2001) [arXiv:hep-th/0007036].
- [24] R. R. Metsaev, C. B. Thorn and A. A. Tseytlin, Nucl. Phys. B **596**, 151 (2001) [arXiv:hep-th/0009171].
- [25] M.B. Green, J. H. Schwarz and L. Brink, Nucl. Phys. **B219**, 437 (1983).
- [26] M.B. Green and J. H. Schwarz, Nucl. Phys. **B243**, 475 (1984); Nucl. Phys. **B218**, 43 (1983).

- [27] J. Maldacena, Adv. Theor. Math. Phys. **2**, 231 (1998) [Int. J. Theor. Phys. **38**, 1113 (1998)] [arXiv:hep-th/9711200].
- [28] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, Phys. Lett. B **428**, 105 (1998) [arXiv:hep-th/9802109].
- [29] E. Witten, Adv. Theor. Math. Phys. **2**, 253 (1998) [arXiv:hep-th/9802150].
- [30] H. Liu and A. A. Tseytlin, Nucl. Phys. B **533**, 88 (1998) [arXiv:hep-th/9804083].
- [31] G. E. Arutyunov and S. A. Frolov, Nucl. Phys. B **544**, 576 (1999) [arXiv:hep-th/9806216].
- [32] R. R. Metsaev, “Light-cone form of field dynamics in AdS space-time,” arXiv:hep-th/0011112. Talk given at DPF 2000: The Meeting of the Division of Particles and Fields of the American Physical Society, Columbus, Ohio, 9-12 Aug 2000.